Brain Research, 490 (1989) 407–409
Elsevier

Author Index

Aijón, J., see Alonso, J.R., 385
Akimoto, K., Hamamura, T. and Otsuki, S., Subchronic cocaine treatment enhances cocaine-induced dopamine efflux, studied by in vivo intracerebral dialysis, 339
Akiyama, A., see Nakazato, T., 332
Akiyama, K., see Yamada, N., 126
Albrecht, J., see Pylova, S.I., 170
Allan, A.M., see Harris, R.A., 26
Alonso, J.R., Coveñas, R., Lara, J., De León, M. and Aijón, J., Distribution of vasoactive intestinal polypeptide-like immunoreactivity in the olfactory bulb of the rainbow trout (*Salmo gairdneri*), 385
Alonso, M.J., see Salaices, M., 133
Anders, J.J., see Hall, L.L., 157
Angel, I. and Taranger, M.-A., Coupling between hypothalamic α_2-adrenoceptors and [^{3}H]mazindol binding site in response to several hyperglycaemic stimuli in mice, 367
Atrens, D.M., see McGregor, I.S., 397

Baker, J.F., see Mason, C.R., 373
Balleine, B.W., see McGregor, I.S., 397
Bambagioni, D., see Grassi, S., 345
Beck, B., see Gaspar, P., 1
Bedoy, C.A. and Mobley, P.L., Astrocyte morphology altered by 1-(5-isoquinolinylsulfonyl) 2-methyl piperazine (H-7) and other protein kinase inhibitors, 243
Benoit, R., see Gaspar, P., 1
Bereiter, D.A. and Gann, D.S., Substance P and GABAergic effects on adrenal and autonomic function evoked by microinjections into trigeminal subnucleus caudalis in the cat, 307
Berger, B., see Gaspar, P., 1
Bernard, J.F., Peschanski, M. and Besson, J.M., Afferents and efferents of the rat cuneiformis nucleus: an anatomical study with reference to pain transmission, 181
Besson, J.M., see Bernard, J.F., 181
Beyer, C., see Pacheco, P., 85
Bird, E.D., see Dodd, P.R., 320
Bondoux Jahan, M. and Sebille, A., Conditioning lesion effects on rat sciatic nerve regeneration are influenced by electrical stimulation delivered to denervated muscles, 350
Borke, R.C., see Hall, L.L., 157
Bowersox, S., see Mignot, E., 186
Brown, L.L., see Justice, A., 73

Cady, A.B., see Obal Jr., F., 292
Ciaranello, R., see Mignot, E., 186
Clascá, F., Llamas, A. and Reinoso-Suárez, F., Hypothalamic connections of the insular cortex in the cat, 361
Claus, H., see Heimrich, B., 404
Cohen, A.I., Interphotoreceptor contacts at the inner segment level in primate retinas, 200
Collins, F. and Crutcher, K.A., Sustained elevation in hippocampal NGF-like biological activity following medial septal lesions in the rat, 355
Coveñas, R., see Alonso, J.R., 385
Cowburn, R.F., see Dodd, P.R., 320
Crutcher, K.A., see Collins, F., 355

DeLanney, L.E., see Ricaurte, G.A., 301
De León, M., see Alonso, J.R., 385
Demediuk, P., see Vink, R., 144
Dement, W.C., see Mignot, E., 186
Dodd, P.R., Watson, W.E.J., Morrison, M.M., Johnston, G.A.R., Bird, E.D., Cowburn, R.F. and Hardy, J.A., Uptake of γ-aminobutyric acid and L-glutamic acid by synaptosomes from postmortem human cerebral cortex: multiple sites, sodium dependence and effect of tissue preparation, 320
Duyckaerts, C., see Gaspar, P., 1

Eilam, D., Golani, I. and Szechtman, H., D_2-agonist quinpirole induces perseveration of routes and hyperactivity but no perseveration of movements, 255
Eimerl, S., see Minc-Golomb, D., 205
Ensor, D.M., Morley, J.S., Redfern, R.M. and Miles, J.B., Stimulation of dopamine output from adrenal medullary cells by β-endorphin and its C-terminal tetrapeptide (MPF), 196

Faden, A.I., see Vink, R., 144
Feasey, K.J., see Stelzer, A., 41
Febvret, A., see Gaspar, P., 1
Feldman, S.M., see Justice, A., 73
Fernández-Alfonso, M.S., see Salaices, M., 133
Fowler, J.C., Adenosine antagonists delay hypoxia-induced depression of neuronal activity in hippocampal brain slice, 378
Frusthofer, B., see Mignot, E., 186
Fujimoto, K., see Ookuma, K., 268
Fukagawa, K., see Ookuma, K., 268

Gann, D.S., see Bereiter, D.A., 307
Gaspar, P., Duyckaerts, C., Febvret, A., Benoit, R., Beck, B. and Berger, B., Subpopulations of somatostatin 28-immunoreactive neurons display different vulnerability in senile dementia of the Alzheimer type, 1
Golani, I., see Eilam, D., 255
Grassi, S., Ottaviani, F. and Bambagioni, D., Vocalization and stapedius muscle activity evoked by local electrical stimulation of midbrain in the chicken (*Gallus gallus*), 345
Gregor, L., see Thornhill, J.A, 174
Guilleminault, C., see Mignot, E., 186

Haas, H.L., see Heimrich, B., 404
Hall, L.L., Borke, R.C. and Anders, J.J., Transection or electrical stimulation of the hypoglossal nerve increases glial fibrillary acidic protein immunoreactivity in the hypoglossal nucleus, 157
Hamamura, T., see Akimoto, K., 339
Harding, J.W., see Jensen, L.L., 48
Hardy, J.A., see Dodd, P.R., 320
Harris, L.R., see Rauschecker, J.P., 56
Harris, R.A. and Allan, A.M., Genetic differences in coupling of benzodiazepine receptors to chloride channels, 26
Heimrich, B., Claus, H., Schwegler, H. and Haas, H.L., Hippocampal mossy fiber distribution and long-term potentiation in two inbred mouse strains, 404

Henegar, J.R., see Maruniak, J.A., 212
Hernandez, L., see Silva, E., 192
Herz, A., see Sirinathsinghji, D.J.S., 276
Hilgier, W., see Pylova, S.I., 170

Irwin, I., see Ricaurte, G.A., 301
Itoh, K., see Ogawa, R., 152

Jensen, L.L., Harding, J.W. and Wright, J.W., Increased blood pressure induced by central application of aminopeptidase inhibitors is angiotensinergic-dependent in normotensive and hypertensive rat strains, 48
Johannsen, L., see Obal Jr., F., 292
Johnston, G.A.R., see Dodd, P.R., 320
Johnston, M.V., see McDonald, J.W., 33
Jones, D.J., see Pangerl, A., 166
Joó, F., Tósaki, A., Oláh, Z. and Koltai, M., Inhibition of H-7 of the protein kinase C prevents formation of brain edema in Sprague–Dawley CFY rats, 141
Justice, A., Feldman, S.M. and Brown, L.L., The nucleus locus coeruleus modulates cerebral glucose utilization during noise stress in rats, 73

Kanazawa, I., see Konishi, S, 162
Kaneko, T., see Ogawa, R., 152
Kapuściński, A., see Pylova, S.I., 170
Kater, S.B., see Mattson, M.P., 110
Kato, H., see Nakano, S., 178
Katz, J.L., see Ricaurte, G.A., 301
Klooster, J., see Ten Tusscher, M.P.M., 95
Kogure, K., see Nakano, S., 178
Koltai, M., see Joó, F., 141
Komisaruk, B.R., see Pacheco, P., 85
Konishi, S., Song, S.-Y., Ogawa, T. and Kanazawa, I., Tachykinins produce fast and slow depolarizations in sympathetic neurons of rat coeliac-superior mesenteric ganglia, 162
Krueger, J.M., see Obal Jr., F., 292

Langston, J.W., see Ricaurte, G.A., 301
Lara, J., see Alonso, J.R., 385
Lemke, M., see Vink, R., 144
Lin, P.J., see Maruniak, J.A., 212
Llamas, A., see Clascá, F., 361

Maddaluno, J., see Mignot, E., 186
Majkowska, J., see Pylova, S.I., 170
Marin, J., see Salaices, M., 133
Martinez-Gomez, M., see Pacheco, P., 85
Maruniak, J.A., Lin, P.J. and Henegar, J.R., Effects of unilateral naris closure on the olfactory epithelia of adult mice, 212
Mason, C.R. and Baker, J.R., Effects of optokinetic velocity and medial vestibulo-cerebellum lesions on vestibulo-ocular reflex direction adaptation in the cat, 373
Mathison, R., see Thornhill, J.A., 174
Mattson, M.P. and Kater, S.B., Development and selective neurodegeneration in cell cultures from different hippocampal regions, 110
McDonald, J.W., Silverstein, F.S. and Johnston, M.V., Neuroprotective effects of MK-801, TCP, PCP and CPP against *N*-methyl-D-aspartate induced neurotoxicity in an in vivo perinatal rat model, 33
McGregor, I.S., Balleine, B.W. and Atrens, D.W., Footshock stress facilitates self-stimulation of the medial prefrontal cortex but not the lateral hypothalamus in the rat, 397
McLaughlin, P.J., see Zagon, I.S., 14
Mignot, E., Guilleminault, C., Bowersox, S., Frusthofer, B., Nishino, S., Maddaluno, J., Ciaranello, R. and Dement, W.C., Central α_1 adrenoceptor subtypes in narcolepsy-cataplexy: a disorder of REM sleep, 186
Miles, J.B., see Ensor, D.M., 196
Minc-Golomb, D., Eimerl, S. and Schramm, M., Cysteine sulfinic acid-induced release of D-[^{3}H]aspartate and [^{14}C]-GABA in hippocampus slices: the role of sodium channels and cAMP, 205
Mizuno, N., see Ogawa, R., 152
Mobley, P.L., see Bedoy, C.A., 243
Moneta, M.E., see Stelzer, A., 41
Mori-Okamoto, J. and Tatsuno, J., Participation of cyclic adenosine monophosphate and β-adrenergic receptors in the facilitatory effect of noradrenaline on the response of cultured cerebellar neurons to glutamate, 64
Moritani, T., see Yamashita, N., 148
Morley, J.S., see Ensor, D.M., 196
Morrison, M.M., see Dodd, P.R., 320

Nakano, S., Kato, H. and Kogure, K., Neuronal damage in the rat hippocampus in a new model of repeated reversible transient cerebral ischemia, 178
Nakazato, T. and Akiyama, A., Effect of exogenous L-DOPA on behavior in the rat: an in vivo voltammetric study, 332
Nazruddin, Suemune, S., Shirana, Y., Yamauchi, K. and Shigenaga, Y., The cells of origin of the hypoglossal afferent nerves and central projections in the cat, 219
Nikolarakis, K.E., see Sirinathsinghji, D.J.S., 276
Nishino, S., see Mignot, E., 186
Noble, E.P., see Stelzer, A., 41

Obal Jr., F., Opp, M., Cady, A.B., Johannsen, L. and Krueger, J.M., Prolactin, vasoactive intestinal peptide, and peptide histidine methionine elicit selective increases in REM sleep in rabbits, 292
Ogawa, R., Itoh, K., Kaneko, T. and Mizuno, N., Co-existence of vasoactive intestinal polypeptide (VIP)- and cholecystokinin (CCK)-like immunoreactivities in thalamocortical neuron in the ventrolateral nucleus of the rat, 152
Ogawa, T., see Konishi, S., 162
Okuno, E., see Schwarcz, R., 103
Oláh, Z., see Joó, F., 141
Ookuma, K., Yoshimatsu, H., Sakata, T., Fujimoto, K. and Fukagawa, K., Hypothalamic sites of neuronal histamine action on food intake by rats, 268
Opp, M., see Obal Jr., F., 292
Otsuki, S., see Akimoto, K., 339
Otsuki, S., see Yamada, N., 126
Ottaviani, F., see Grassi, S., 345

Pacheco, P., Martinez-Gomez, M., Whipple, B., Beyer, C. and Komisaruk, B.R., Somato-motor components of the pelvic and pudendal nerves of the female rat, 85
Pangerl, A., Pangerl, B., Reiter, R.J., Vaughan, G.M. and Jones, D.J., Twenty-four hour variation of α_1-adrenergic receptors in the pineal gland of the male Syrian hamster, 166
Pangerl, B., see Pangerl, A., 166
Peschanski, M., see Bernard, J.F., 181
Pittman, Q., see Thornhill, J.A., 174
Pylova, S.I., Majkowska, J., Hilgier, W., Kapuściński, A. and Albrecht, J., Rapid decrease of high affinity ouabain binding

sites in hippocampal CA1 regions following short-term global cerebral ischemia in rat, 170

Rauschecker, J.P. and Harris, L.R., Auditory and visual neurons in the cat's superior colliculus selective for the direction of apparent motion stimuli, 56
Redfern, R.M., see Ensor, D.M., 196
Reinoso-Suárez, F., see Clascá, F., 361
Reiter, R.J., see Pangerl, A., 166
Ricaurte, G.A., DeLanney, L.E., Irwin, I., Witkin, J.M., Katz, J.L. and Langston, J.W., Evaluation of the neurotoxic potential of *N,N*-dimethylamphetamine: an illicit analog of methamphetamine, 301
Rico, I., see Salaices, M., 133

Sakata, T., see Ookuma, K., 268
Salaices, M., Alonso, M.J., Rico, I., Fernández-Alfonso, M.S. and Marin, J., Vasoconstrictor effect of the Ca^{2+} agonist Bay K 8644 on human cerebral arteries, 133
Schramm, M., see Minc-Golomb, D., 205
Schwarcz, R., Okuno, E. and White, R.J., Basal ganglia lesions in the rat: effects on quinolinic acid metabolism, 103
Schwegler, H., see Heimrich, B., 404
Schweitzer, J.B., Nerve growth factor receptor-mediated transport from CSF labels cholinergic neurons: direct demonstration by a double-labeling study, 390
Sebille, A., see Bondoux-Jahan, M., 350
Shigenaga, Y., see Nazruddin, 219
Shirana, Y., see Nazruddin, 219
Silva, E. and Hernandez, L., Goldthioglucose causes brain norepinephrine and serotonin depletion correlated with increased body weight, 192
Silverstein, F.S., see McDonald, J.W., 33
Simson, P.E. and Weiss, J.M., Peripheral, but not local or intracerebroventricular, administration of benzodiazepines attenuates evoked activity of locus coeruleus neurons, 236
Sincini, E., see Stelzer, A., 41
Sirinathsinghji, D.J.S., Nikolarakis, K.E. and Herz, A., Corticotropin-releasing factor stimulates the release of methionine-enkephalin and dynorphin from the neostriatum and globus pallidus of the rat: in vitro and in vivo studies, 276
Song, S.-Y., see Konishi, S., 162
Stelzer, A., Feasey, K.J., Moneta, M.E., Sincini, E., Ten Bruggencate, G. and Noble, E.P., Inositol 1-phosphate formation in long-term potentiation and kindling, 41
Suemune, S., see Nazruddin, 219
Szechtman, H., see Eilam, D., 255

Taranger, M.-A., see Angel, I., 367
Tatsuno, J., see Mori-Okamoto, J., 64
Ten Bruggencate, G., see Stelzer, A., 41
Ten Tusscher, M.P.M., Klooster, J. and Vrensen, G.F.J.M., Satellite cells as blood–ganglion cell barrier in autonomic ganglia, 95
Thornhill, J.A., Gregor, L., Mathison, R. and Pittman, Q., Intrathecal dynorphin A administration causes pressor responses in rats associated with an increased resistance to spinal cord blood flow, 174
Tósaki, A., see Joó, F., 141

Vaughan, G.M., see Pangerl, A., 166
Vink, R., Yum, S.W., Lemke, M., Demediuk, P. and Faden, A.I., Traumatic spinal cord injury in rabbits decreases intracellular free magnesium concentration as measured by ^{31}P MRS, 144
Vrensen, G.F.J.M., see Ten Tusscher, M.P.M., 95

Watson, W.E.J., see Dodd, P.R., 320
Weiss, J.M., see Simson, P.E., 236
Whipple, B., see Pacheco, P., 85
White, R.J., see Schwarcz, R., 103
Witkin, J.M., see Ricaurte, G.A., 301
Wright, J.W., see Jensen, L.L., 48

Yamada, N., Akiyama, K. and Otsuki, S., Hippocampal kindling enhances excitatory amino acid receptor-mediated polyhosphoinositide hydrolysis in the hippocampus and amygdala/pyriform cortex, 126
Yamashita, N. and Moritani, T., Anticipatory changes of soleus H-reflex amplitude during execution process for heel raise from standing position, 148
Yamauchi, K., see Nazruddin, 219
Yoshimatsu, H., see Ookuma, K., 268
Yum, S.W., see Vink, R., 144

Zagon, I.S. and McLaughlin, P.J., Endogenous opioid systems regulate growth of neural tumor cells in culture, 14